



Impact of obesity on incident hypertension independent of weight gain among nonhypertensive Japanese: the Ibaraki Prefectural Health Study (IPHS)

著者	Tsujimoto Takehiko, Sairenchi Toshimi, Iso Hiroyasu, Irie Fujiko, Yamagishi Kazumasa, Tanaka Kiyoji, Muto Takashi, Ota Hitoshi
journal or publication title	Journal of hypertension
volume	30
number	6
page range	1122-1128
year	2012-06
権利	(C) 2012 Wolters Kluwer Health Lippincott Williams & Wilkins. This is a non-final version of an article published in final form in Journal of Hypertension: June 2012 - Volume 30 - Issue 6 - p 1122 1128, doi: 10.1097/HJH.0b013e328352b879
URL	http://hdl.handle.net/2241/119507

doi: 10.1097/HJH.0b013e328352b879

Title page

Title

Impact of Obesity on Incident Hypertension Independent of Weight Gain among
Non-hypertensive Japanese: The Ibaraki Prefectural Health Study (IPHS)

Authors

Takehiko TSUJIMOTO (MS)^{a, b}, Toshimi SAIRENCHI (Ph.D.)^{a, c}, Hiroyasu ISO (MD, Ph.D.,
MPH)^d, Fujiko IRIE (MD, Ph.D.)^e, Kazumasa YAMAGISHI (MD, Ph.D.)^{a, b}, Kiyoji
TANAKA (Ph.D.)^b, Takashi MUTO (MD, Ph.D.)^c, Hitoshi OTA (MD, Ph.D.)^a

a. Ibaraki Health Plaza, Ibaraki Health Service Association, Ibaraki, Japan

b. Graduate School of Comprehensive Human Sciences, University of Tsukuba, Ibaraki,
Japan

c. Department of Public Health, Dokkyo Medical University School of Medicine, Tochigi,
Japan

d. Department of Social and Environmental Medicine, Graduate School of Medicine,
Osaka University, Osaka, Japan

e. Department of Health and Welfare, Ibaraki Prefectural Office, Ibaraki, Japan

Running title: Stable obesity and incident hypertension

Word counts: 2 720 (text), 248 (abstract), 2 tables, 1 figure, 35 references

Corresponding author

Toshimi SAIRENCHI, Ph.D.

Department of Public Health, Dokkyo Medical University School of Medicine

Adress: 880 Kita-kobayashi, Mibu, Tochigi 321-0293, Japan

Phone: +81-282-87-2133, Fax: +81-282-86-2935

E-mail: tossair@dokkyomed.ac.jp

Abstract

Objective: The aim of this study was to examine the association between body mass index (BMI) and risk of incident hypertension among Japanese men and women who are middle-aged and older.

Design: Prospective, population-based cohort study.

Subjects: A total of 68 205 non-hypertensive adults (18 336 men and 49 869 women) aged 40-79 years who completed health check-ups in the Ibaraki prefecture, Japan, in 1993 were followed-up through 2006. To exclude the impact of BMI change during the follow-up period, a time-dependent covariate Cox proportional hazards model was used to compute the hazard ratios (HRs) of incident hypertension according to BMI categories. Incident hypertension was defined as a systolic blood pressure (BP) of ≥ 140 mmHg, a diastolic BP of ≥ 90 mmHg, and/or hypertensive medication use.

Results: A total of 30 982 adults (45.4%) developed hypertension (9 331 men and 21 651 women) during a mean of 3.9 years of follow-up. Compared with a BMI of < 19.0 , time-dependent covariates adjusted HRs (95% confidence interval [CI]) for hypertension among participants with a BMI of ≥ 25.0 were 1.42 (1.17-1.73) for men aged 40 to 59 years, 1.34 (1.19-1.51) for men aged 60 to 79 years, 1.47 (1.33-1.62) for women aged 40 to 59 years, and 1.29 (1.18-1.41) for women aged 60 to 79 years.

Conclusion: The baseline BMI is associated with future risk for incident hypertension even after accounting for weight change during the follow-up period. Weight loss may be recommended to non-hypertensive obese adults to prevent the development of hypertension.

Introduction

Hypertension is a major cause of cardiovascular disease [1]. Currently, approximately 45% of Japanese adults are considered hypertensive [2], defined as systolic blood pressure (SBP) \geq 140 mmHg, diastolic blood pressure (DBP) \geq 90 mmHg or treatment with an antihypertensive drug. Many prospective cohort studies have consistently shown that the higher the BMI, the greater the likelihood of developing hypertension [3-11]. Some prospective cohort studies have reported that long-term weight gain was associated with a risk of hypertension among hypotensive or high-normal blood pressure patients [12-15]. Moreover, many intervention studies have also shown that weight loss reduced blood pressure [16-18]. These results could lead to a hypothesis that the association between baseline BMI and the risk of hypertension are due to weight gain during the follow-up period. If this hypothesis were true, non-hypertensive obese individuals would not be required to lose weight to prevent incident hypertension. To clarify whether weight loss should be recommended for non-hypertensive obese people to prevent incident hypertension, a large cohort study that accounts for weight change is warranted. The purpose of this study was to clarify the association between long-term weight stability and the risk of incident hypertension in a large community-based cohort of men and women who are middle-aged and older.

Methods

Study population

The study population comprised 194 333 individuals (63 865 men and 130 468 women) aged 40-79 years living in the Ibaraki Prefecture, Japan. The population participated in community-based annual health checkups beginning in 1993 (Ibaraki Prefectural Health Study). These health checkups were conducted by local governments in accordance with the Japan Health Laws. Data were collected by the Ibaraki prefectural government from local governments after depersonalizing them to ensure anonymity. We excluded 3 080 individuals (738 men and 2 342 women) with incomplete data and 94 153 individuals (35 647 men and 58 506 women) with a SBP of ≥ 140 mmHg and/or a DBP of ≥ 90 mmHg or antihypertensive medication use. We further excluded 28 895 individuals (9 144 men and 19 751 women) who did not participate in the 1994 survey, thereby ensuring that the participants were followed for at least one year.

Ultimately, the study subjects consisted of 68 205 individuals (18 336 men and 49 869 women). These participants were followed up by annual examinations until hypertension had been diagnosed or until the end of 2006. Blood pressure was measured at annual follow-up examinations. Participants who did not undergo annual checkups during the follow-up periods were censored on the date of their latest checkup. The protocol for this cohort study was approved by the Ibaraki Epidemiology Study Union Ethics Review Committee.

Assessment of body mass index, blood pressure, and other covariates

Height in stockinged feet and weight in light clothing were measured at baseline. BMI was calculated as weight in kilograms divided by the height in meters squared. BP was measured from the right arm of seated participants who had rested for more than 5 minutes by trained observers using standard mercury sphygmomanometers during examinations that occurred

between 1993 and 2004 and by an automated sphygmomanometer between 2005 and 2006. When the SBP was greater than 150 mmHg or DBP was greater than 90 mmHg, a second measurement was performed after several deep breaths, and the lower values, which were almost always observed after the second measurement, were used for analyses.

Blood samples were drawn from seated participants into two polyethylene terephthalate tubes: one with an accelerator and the other with sodium fluoride and ethylenediaminetetraacetic acid. Overnight fasting (≥ 8 hours) was not required. The blood glucose level was measured by the glucose oxidase electrode method with a GA1140 device (Kyoto Daiichi Kagaku, Kyoto, Japan) during examinations that occurred between 1993 and 1996, by the enzyme method with a H7170 device (Hitachi, Tokyo, Japan) between 1997 and 2003, and with a H7700 device (Hitachi, Tokyo, Japan) between 1994 and 2006. Participants were considered diabetic if they had a fasting plasma glucose level of at least 6.1 mmol/L or a non-fasting glucose level of at least 7.8 mmol/L, or if they were being treated for diabetes mellitus. Serum total cholesterol and serum triglyceride levels were measured by the enzyme method with a RX-30 device (Nihon Denshi, Tokyo, Japan) between 1993 and 1995, with a H7350 device (Hitachi, Tokyo, Japan) between 1996 and 2003, and with a H7700 device between 2004 and 2006. High-density lipoprotein (HDL) cholesterol levels were measured by the phosphotungstic acid magnesium method with a MTP-32 device (Corona Electric, Ibaraki, Japan) between 1993 and 1995, by the selective inhibition method with a H7350 device between 1996 and 2003, and with a H7700 device between 2004 and 2006. The laboratory participated in external standardization and successfully met the criteria for precision accuracy for the measurement of blood samples by the Japan Medical Association, the Japanese Association of Medical Technologists, and the Japan Society of Health Evaluation and Promotion. An interview was conducted to ascertain medical history, smoking status (never, ex-, current < 20 cigarettes/day, and ≥ 20 cigarettes/day), and alcohol intake (non-,

sometimes, < 66 g/day, and \geq 66 g/day).

Endpoint determination

Incident hypertension was defined as SBP of \geq 140 mmHg, DBP of \geq 90 mmHg, and/or when treatment for hypertension was initiated.

Statistical analysis

Participants were classified into the following 7 categories with regard to their BMI: < 19.0, 19.0-20.9, 21.0-22.9, 23.0-24.9, 25.0-26.9, 27.0-29.9, or \geq 30.0 kg/m². Baseline participant characteristics were compared according to BMI categories using chi-squared tests for categorical variables and analysis of variance for continuous variables. Hazard ratios (HRs) with 95% confidence intervals (CIs) for incident hypertension relative to BMI categories were calculated with a reference of < 19.0 kg/m² using a Cox proportional hazards regression model [19]. The analyses were stratified by sex and age groups (40-59 and 60-79 years).

Three multivariable-adjusted models were used. In Model 1, covariates included age (years), SBP (mmHg), DBP (mmHg), fasting status (yes or no), total cholesterol level (mmol/L), HDL cholesterol level (mmol/L), log-transformed triglyceride level (mmol/L), lipid medication use (yes or no), blood glucose status (normal: < 6.1 mmol/L during fasting or < 7.8 mmol/L during non-fasting; border: 6.1-7.0 mmol/L during fasting or 7.8-11.1 mmol/L during non-fasting; hyperglycemia: \geq 7.0 mmol/L during fasting or \geq 11.1 mmol/L during non-fasting), smoking status (never, ex-, current < 20 cigarettes/day, or \geq 20 cigarettes/day), and alcohol intake (non-, sometimes, < 66 g/day, or \geq 66 g/day). In Model 2, the changes in BMI, total cholesterol level (mmol/L), HDL cholesterol level (mmol/L), and log-transformed triglyceride level (mmol/L) from baseline to the end of follow-up were added as covariates to Model 1. In Model 3, time-dependent variables of BMI and covariates

in Model 1 were used with a time-dependent Cox hazard model, which was used for adjusting the change in the variable from the baseline to the final year of follow-up [20]. For the secondary analysis, we excluded 34 190 participants with incomplete data and 16,370 participants with a SBP of ≥ 140 mmHg and/or a DBP of ≥ 90 mmHg or antihypertensive medication use at a 5-year follow-up. The remaining 17 645 qualified participants were classified into the following 4 categories with regard to their baseline BMI and BMI 5 years later: (a) BMI of less than 25 kg/m^2 at baseline and BMI of less than 25 kg/m^2 at a 5-year follow-up (-/-); (b) BMI of 25 kg/m^2 or more at baseline and BMI of less than 25 kg/m^2 at a 5-year follow-up (+/-); (c) BMI of less than 25 kg/m^2 at baseline and BMI of 25 kg/m^2 or more at a 5-year follow-up (-/+); and (d) BMI of 25 kg/m^2 or more at baseline and BMI of 25 kg/m^2 or more at a 5-year follow-up (+/+). We calculated HRs for incident hypertension to consider long-term changes of their weights.

All statistical analyses were conducted using SAS, version 9.1 (SAS Institute, Inc., Cary, North Carolina).

Results

Sex-stratified baseline characteristics according to BMI categories are shown in Table 1. All covariates, except for blood glucose status, diabetic medication use and smoking status in men and lipid medication use in women, were associated with BMI in both sexes. A higher BMI was linked with higher SBP and DBP at baseline in both sexes.

Of the 68 205 adults (18 336 men and 49 869 women), 30 982 (45.4%) developed hypertension (9 331 men and 21 651 women) during a mean follow-up of 3.9 years (3.4 years for men and 4.1 years for women). Table 2 shows sex- and age-stratified HRs for hypertension according to the baseline BMI categories. In men aged 40-59 years, compared with a baseline BMI $< 19.0 \text{ kg/m}^2$, the multivariate HRs of BMI $\geq 25.0 \text{ kg/m}^2$ was significantly higher in Model 1 and Model 2. In men aged 60-79 years, compared with a baseline BMI $< 19.0 \text{ kg/m}^2$, the multivariate HRs of BMI $\geq 19.0 \text{ kg/m}^2$ was significantly higher in Model 1 and the multivariate HRs of BMI $\geq 23.0 \text{ kg/m}^2$ was significantly higher in Model 2. In both age groups among men, time-dependent covariates adjustment HRs (BMI $\geq 25.0 \text{ kg/m}^2$: 1.42, $\geq 30.0 \text{ kg/m}^2$: 1.77 in aged 40-59 years; BMI $\geq 25.0 \text{ kg/m}^2$: 1.34, $\geq 30.0 \text{ kg/m}^2$: 1.68 in aged 60-79 years) showed a trend similar to Model 1 and Model 2.

In women aged 40-59 years, compared with a baseline BMI $< 19.0 \text{ kg/m}^2$, the multivariate HRs of BMI $\geq 21.0 \text{ kg/m}^2$ was significantly higher in Model 1 and of BMI $\geq 23.0 \text{ kg/m}^2$ was significantly higher in Model 2.

In women aged 60-79 years, compared with a baseline BMI $< 19.0 \text{ kg/m}^2$, the multivariate HRs of BMI $\geq 21.0 \text{ kg/m}^2$ was significantly higher in Model 1 and 25.0-26.9 kg/m^2 of BMI was significantly higher in Model 2. In both age groups among women, time-dependent covariates adjustment HRs (BMI $\geq 25.0 \text{ kg/m}^2$: 1.53, $\geq 30.0 \text{ kg/m}^2$: 2.07 in age 40-59 years; BMI $\geq 25.0 \text{ kg/m}^2$: 1.47, $\geq 30.0 \text{ kg/m}^2$: 1.91 in age 60-79 years) showed a trend similar to Model 1 and Model 2.

Figure 1 represents HRs for hypertension according to the BMI categories at baseline and at a 5-year follow-up. Among all non-hypertensive participants at baseline and at a 5-year follow-up, compared with the -/- group, the age-adjusted HRs (95% CI) of +/-, -/+, and +/+ were 1.12 (1.03-1.41), 1.69 (1.47-1.94), and 1.67 (1.53-1.82), respectively. The multivariate HRs of +/-, -/+, and +/+ were 1.12 (1.03-1.41), 1.69 (1.47-1.94), and 1.67 (1.53-1.82), respectively.

Discussion

To the best of our knowledge, the results of the present large prospective study showed that stable obesity in non-hypertensive individuals was significantly associated with an increased risk of incident hypertension among men and women who are middle-aged and older.

Previous studies that examined the relationship between BMI and hypertension reported results consistent with the present study. Many cross-sectional studies showed a dose-dependent relationship between BMI and the prevalence of hypertension in Caucasian [5, 21] and Asian [3, 13, 22] populations. Kawada [3] reported that the odds ratios for hypertension against the lowest quartile of BMI (separated by 20, 25 and 28) were 1.7, 3.6, and 7.2, respectively, with 95% CIs of 1.2-2.3, 2.5-5.2 and 4.7-11.0, respectively, among middle-aged men and women (5 346 men and 11 525 women, aged 40-59). Prospective studies [4, 12, 23] also showed similar results to the present study. The Physicians' Health Study (13 563 healthy and non-hypertensive men in U.S.) [4] showed that the HRs (95% CIs) of developing hypertension for men with a BMI of 22.4-23.6, 23.7-24.7, 24.8-26.4, and >26.4 kg/m² were 1.20 (1.09-1.32), 1.31 (1.19-1.44), 1.56 (1.42-1.72), and 1.85 (1.69-2.03), respectively (*P* for the trend, < 0.001), compared with participants who had the lowest BMI quintile (< 22.4 kg/m²). A cohort study of U.S. female nurses (82 473 women, aged 30-55) [12] showed that the multivariate HRs (95% CIs) of developing hypertension for women with a BMI of 22.0-22.9, 24.0-24.9, 26.0-27.9, and > 31.0 kg/m² were 1.57 (1.44-1.72), 2.15 (1.97-2.35), 3.33 (3.06-3.62), and 6.31 (5.80-6.87), respectively (*P* for the trend, < 0.001), compared with women who had the lowest BMI decile (< 20.0 kg/m²), and an increase in BMI of 1 kg/m² was associated with a 12 % increased risk for incident hypertension. These results suggested a strong association between higher BMI and increased risk of incident hypertension in men and women.

We also examined the effect of 5-year changes in BMI category (BMI less than, or equal or

more than 25 kg/m²) on incident hypertension. The HRs of non-obese people at a 5-years follow-up, despite obesity at baseline, was not significant compared with non-obese people at both baseline and at a 5-years follow-up. In contrast, the HRs of obese individuals at a 5-year follow-up, despite non-obesity at baseline, was significantly higher compared with non-obese people at both baseline and at a 5-years follow-up. The effect of change in BMI on blood pressure has already been examined in some studies [12-18]. A systematic review [24] of lifestyle intervention studies reported differences ranging from -11 to + 4 kg for weight, -7 to + 2.2 mmHg for DBP and -13 to + 6.1 mmHg for SBP. It is suggested that SBP differences were equal to a ratio of 1 kg to every 1 mmHg. These studies suggest the efficacy of weight loss for the hypertensive obese population. The present study further represented the efficacy of weight loss for the non-hypertensive obese population to prevent incident hypertension. Williams [25] examined the relationship between long-term stable individuals' BMI (change less than ± 0.4 kg/m²) and the cumulative incidence of hypertension and showed that the odds ratio of each kg/m² increment in baseline BMI was 1.19 (95% CIs: 1.14-1.24) in men and 1.11 (95% CIs: 1.02-1.20) in women. Matsuo et al [26] reported in a cohort study of 5 201 Japanese middle-aged male employees using of a time-dependent Cox proportional hazard model for analysis that the HRs (95% CIs) of stable BMI of 25.0-26.9, 27.0-29.9, and ≥ 30.0 kg/m² were 1.11 (0.93-1.32), 1.34 (1.09-1.65) and 1.49 (1.09-2.04), respectively, compared with men who had a BMI of 23.0-24.9 kg/m². These results are consistent with our results, although Williams calculated the BMI with a self-reported weight and cumulative incidence of hypertension at 2 points (baseline and 7 years later), and Matsuo et al. studied middle-aged male employees.

The exact mechanism by which stable adiposity raises BP is uncertain. However, a likely explanation is that obesity-associated hypertension appears to be due to inadequate vasodilatation in the setting of increased blood volume and cardiac output. A defect in control

of vascular resistance has been attributed to enhanced activity of the sympathetic nervous system, abnormal rennin-angiotensin-aldosterone relations, and insulin resistance [27-31]. The peripheral vascular resistance is increased by adipocytokines (e.g., leptin, TNF- α , RBP4) [32-34].

The study has several strengths. First, the effect of weight change during the follow-up period was taken into account to investigate the association between long-term weight stability and future incident risk for hypertension, while many previous epidemiological studies did not account for these factors. Second, the present study had a large cohort for which gender, age and BMI stratification analysis was possible. In addition, all blood samples in each year were measured by the same device, reagents, and quality control programs [35].

However, there are several limitations. First, the participation rate for the follow-up examinations was approximately 40%, but the mean SBP and DBP did not differ between individuals who were and were not followed up. In this context, the potential selection bias may have been small. Second, the instrument used to measure the blood pressure was switched from the standard mercury sphygmomanometer to an automatic sphygmomanometer following 2005. The BP determined by means of an automated sphygmomanometer method was compared with those determined by a standard mercury sphygmomanometer method, using 18 859 samples (non-medication participants) in 2004 and 2005. Comparability between BP measured using the 2 sphygmomanometers was acceptable (SBP: $r = 0.69$, DBP: $r = 0.55$, $P < 0.05$, respectively). Third, we could not assess the potential confounding variables such as fat distribution, physical activity, nutritional status (e.g., sodium intake), and family history of hypertension.

In conclusion, the baseline BMI is associated with a future incident risk for hypertension, even after accounting for weight change during the follow-up period. Weight loss may be recommended for obese non-hypertensive individuals to prevention incident hypertension

267 among men and women who are middle-aged and older.

268

269 **Acknowledgement**

270 We thank all the participants in this study.

271

272 **Conflict of interest**

273 The authors declare no conflict of interest.

274

References

1. Kannel WB. Role of blood pressure in cardiovascular disease: the Framingham Study. *Angiology* 1975; 26: 1-14.
2. Japanese Ministry of Health and Welfare. National Health and Nutrition Survey. In: Tokyo, Japan: Japanese Ministry of Health and Welfare, 2006. (<http://www.mhlw.go.jp/bunya/kenkou/eiyou08/dl/01-03.pdf>). (Accessed March 20, 2011).
3. Kawada T. Body mass index is a good predictor of hypertension and hyperlipidemia in a rural Japanese population. *Int J Obes Relat Metab Disord* 2002; 26: 725-729.
4. Gelber RP, Gaziano JM, Manson JE, Buring JE, Sesso HD. A prospective study of body mass index and the risk of developing hypertension in men. *Am J Hypertens* 2007; 20: 370-377.
5. Williams PT, Hoffman K, La I. Weight-related increases in hypertension, hypercholesterolemia, and diabetes risk in normal weight male and female runners. *Arterioscler Thromb Vasc Biol* 2007; 27: 1811-1819.
6. Nemesure B, Wu SY, Hennis A, Leske MC. The relationship of body mass index and waist-hip ratio on the 9-year incidence of diabetes and hypertension in a predominantly African-origin population. *Ann Epidemiol* 2008; 18: 657-663.
7. Nguyen TT, Adair LS, He K, Popkin BM. Optimal cutoff values for overweight: using body mass index to predict incidence of hypertension in 18- to 65-year-old Chinese adults. *J Nutr* 2008; 138: 1377-1382.
8. Pang W, Sun Z, Zheng L, Li J, Zhang X, Liu S et al. Body mass index and the prevalence of prehypertension and hypertension in a Chinese rural population. *Intern Med* 2008; 47: 893-897.
9. Shuger SL, Sui X, Church TS, Meriwether RA, Blair SN. Body mass index as a

- predictor of hypertension incidence among initially healthy normotensive women. *Am J Hypertens* 2008; 21: 613-619.
10. Willett WC, Dietz WH, Colditz GA. Guidelines for healthy weight. *N Engl J Med* 1999; 341: 427-434.
11. Chei CL, Iso H, Yamagishi K, Tanigawa T, Cui R, Imano H et al. Body fat distribution and the risk of hypertension and diabetes among Japanese men and women. *Hypertens Res* 2008; 31: 851-857.
12. Huang Z, Willett WC, Manson JE, Rosner B, Stampfer MJ, Speizer FE et al. Body weight, weight change, and risk for hypertension in women. *Ann Intern Med* 1998; 128: 81-88.
13. Ishikawa-Takata K, Ohta T, Moritaki K, Gotou T, Inoue S. Obesity, weight change and risks for hypertension, diabetes and hypercholesterolemia in Japanese men. *Eur J Clin Nutr* 2002; 56: 601-607.
14. Yang G, Xiang YB, Zheng W, Xu WH, Zhang X, Li HL et al. Body weight and weight change in relation to blood pressure in normotensive men. *J Hum Hypertens* 2007; 21: 45-52.
15. Chen PC, Sung FC, Su TC, Chien KL, Hsu HC, Lee YT. Two-year change in body mass index and subsequent risk of hypertension among men and women in a Taiwan community. *J Hypertens* 2009; 27: 1370-1376.
16. He J, Whelton PK, Appel LJ, Charleston J, Klag MJ. Long-term effects of weight loss and dietary sodium reduction on incidence of hypertension. *Hypertension* 2000; 35: 544-549.
17. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA* 2003; 289: 2560-2572.

18. Moore LL, VISIONI AJ, Qureshi MM, Bradlee ML, Ellison RC, D'Agostino R. Weight loss in overweight adults and the long-term risk of hypertension: the Framingham study. *Arch Intern Med* 2005; 165: 1298-1303.
19. Cox DR, Oakes D. *Analysis of Survival Data*. Chapman and Hall, New York 1984.
20. Fisher LD, Lin DY. Time-dependent covariates in the Cox proportional-hazards regression model. *Annu Rev Public Health* 1999; 20: 145-157.
21. Bays HE, Chapman RH, Grandy S. The relationship of body mass index to diabetes mellitus, hypertension and dyslipidaemia: comparison of data from two national surveys. *Int J Clin Pract* 2007; 61: 737-747.
22. Liu L, Choudhury SR, Okayama A, Hayakawa T, Kita Y, Ueshima H. Changes in body mass index and its relationships to other cardiovascular risk factors among Japanese population: results from the 1980 and 1990 national cardiovascular surveys in Japan. *J Epidemiol* 1999; 9: 163-174.
23. Hu G, Barengo NC, Tuomilehto J, Lakka TA, Nissinen A, Jousilahti P. Relationship of physical activity and body mass index to the risk of hypertension: a prospective study in Finland. *Hypertension* 2004; 43: 25-30.
24. Aucott L, Rothnie H, McIntyre L, Thapa M, Waweru C, Gray D. Long-term weight loss from lifestyle intervention benefits blood pressure?: a systematic review. *Hypertension* 2009; 54: 756-762.
25. Williams PT. Increases in weight and body size increase the odds for hypertension during 7 years of follow-up. *Obesity (Silver Spring)* 2008; 16: 2541-2548.
26. Matsuo T, Sairenchi T, Suzuki K, Tanaka K, Muto T. Long-term stable obesity increases risk of hypertension. *Int J Obes (Lond)* 2011; 35: 1056-1062.
27. Julius S, Valentini M, Palatini P. Overweight and hypertension: a 2-way street? *Hypertension* 2000; 35: 807-813.

28. Esler M, Rumantir M, Wiesner G, Kaye D, Hastings J, Lambert G. Sympathetic nervous system and insulin resistance: from obesity to diabetes. *Am J Hypertens* 2001; 14: 304S-309S.
29. Hall JE, Hildebrandt DA, Kuo J. Obesity hypertension: role of leptin and sympathetic nervous system. *Am J Hypertens* 2001; 14: 103S-115S.
30. Verdecchia P, Angeli F. Natural history of hypertension subtypes. *Circulation* 2005; 111: 1094-1096.
31. Poirier P, Giles TD, Bray GA, Hong Y, Stern JS, Pi-Sunyer FX et al. Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss. *Arterioscler Thromb Vasc Biol* 2006; 26: 968-976.
32. Dustan HP. Obesity and hypertension in blacks. *Cardiovasc Drugs Ther* 4 Suppl 1990; 2: 395-402.
33. Patel JV, Lim HS, Dubb K, Hughes EA, Lip GY. Circulating levels of adiponectin, leptin, and tumour necrosis factor alpha in hypertension. *Ann Med* 2009; 41: 291-300.
34. Chiba M, Saitoh S, Ohnishi H, Akasaka H, Mitsumata K, Furukawa T et al. Associations of metabolic factors, especially serum retinol-binding protein 4 (RBP4), with blood pressure in Japanese--the Tanno and Sobetsu study. *Endocr J* 2010; 57: 811-817.
35. Nakamura M, Sato S, Shimamoto T. Improvement in Japanese clinical laboratory measurements of total cholesterol and HDL-cholesterol by the US Cholesterol Reference Method Laboratory Network. *J Atheroscler Thromb* 2003; 10: 145-153.

Figure legend

Figure 1

Abbreviations: HRs, hazard ratios; CIs, confidence intervals. Age-adjusted HRs with 95% confidence intervals were adjusted for age (years) in 1998 and sex. Multivariate HRs with 95% confidence intervals were adjusted for possible confounders measured in 1998: age (years), sex, systolic blood pressure (mmHg), diastolic blood pressure (mmHg), fasting status (yes or no), total cholesterol level (mmol/liter), high density lipoprotein cholesterol level (mmol/liter), log-transformed triglyceride level (mmol/liter), lipid medication use (yes or no), blood glucose status (normal, border, hyperglycemia), diabetic medication use (yes or no), smoking status (never, ex-, < 20 cigarettes/day, or \geq 20 cigarettes/day), and alcohol intake (never, sometimes, < 66 g/day, or \geq 66 g/day). Bold values showed statistical significance ($P < 0.05$).

Table 1. Baseline characteristics of participants by body mass index (BMI) categories

Gender and baseline variables	Body mass index, kg/m ²							<i>P</i> -value
	< 19.0	19.0 - 20.9	21.0 - 22.9	23.0 - 24.9	25.0 - 26.9	27.0 - 29.9	> 30.0	
Men (n = 18 336)								
No. of subjects	1 545	3 476	4 934	4 544	2 598	1 085	154	
Age, years	63.5 (9.5)	60.5 (10.2)	58.8 (10.3)	57.7 (10.2)	56.5 (10.0)	56.6 (9.8)	55.3 (10.0)	<.001
Change in body mass index, kg/m ²	0.0 (1.1)	0.1 (1.2)	0.0 (1.2)	-0.1 (1.3)	-0.2 (1.3)	-0.4 (1.5)	-0.7 (1.5)	<.001
Systolic blood pressure, mmHg	119.9 (11.4)	121.7 (10.8)	122.7 (10.2)	124.0 (9.5)	124.4 (9.4)	125.4 (9.2)	126.2 (8.8)	<.001
Diastolic blood pressure, mmHg	71.9 (8.4)	73.1 (8.0)	74.1 (7.9)	75.4 (7.3)	75.9 (7.4)	76.6 (7.1)	78.1 (7.8)	<.001
Total cholesterol, mmol/liter	4.7 (0.8)	4.8 (0.8)	5.0 (0.8)	5.1 (0.9)	5.2 (0.8)	5.2 (0.9)	5.2 (0.9)	<.001
HDL cholesterol, mmol/liter	1.6 (0.4)	1.5 (0.4)	1.4 (0.4)	1.3 (0.3)	1.2 (0.3)	1.1 (0.3)	1.1 (0.3)	<.001
Triacylglycerol, mmol/liter	1.1 (0.5)	1.2 (0.7)	1.5 (0.8)	1.8 (1.0)	2.0 (1.2)	2.2 (1.3)	2.2 (1.0)	<.001
Lipid medication use, %	3.2	2.8	2.6	3.1	3.2	3.1	2.6	<.001
Diabetic medication use, %	0.4	0.8	1.0	1.3	1.2	1.4	2.6	0.177
Blood glucose status, %								0.866
Normal	81.9	83.9	84.3	84.6	81.9	82.0	83.8	
Border	13.6	12.4	12.0	11.4	13.5	12.7	9.7	
Hyperglycemia	4.5	3.7	3.7	4.0	4.5	5.3	6.5	
Smoking status, %								0.116
Never	19.4	20.6	22.4	25.1	24.1	23.3	25.3	
Ex	23.4	23.3	27.0	28.1	29.5	33.9	27.9	
Current								
< 20 cigarettes/day	22.8	18.7	15.3	12.3	11.5	9.8	9.7	
≥ 20 cigarettes/day	34.4	37.4	35.3	34.4	34.9	33.0	37.0	
Alcohol intake, %								<.001
Never	48.5	40.6	37.2	36.6	37.6	38.2	51.3	
Sometimes	11.5	13.2	13.6	15.1	15.9	18.8	17.5	
Everyday								
< 66 g/day	36.6	41.9	44.0	43.4	41.4	37.1	25.3	
≥ 66 g/day	3.4	4.3	5.1	5.0	5.1	5.8	5.8	

Table 1. Baseline characteristics of participants by body mass index (BMI) categories (continued)

Gender and baseline variables	Body mass index, kg/m ²								P-value
	< 19.0	19.0 - 20.9	21.0 - 22.9	23.0 - 24.9	25.0 - 26.9	27.0 - 29.9	> 30.0		
Women (n = 49 869)									
No. of subjects	3 796	9 391	13 824	11 887	6 781	3 434	756		
Age, years	56.8 (10.9)	54.1 (10.1)	54.4 (9.7)	55.1 (9.5)	56.0 (9.4)	56.1 (9.4)	55.4 (9.2)		<.001
Change in body mass index, kg/m ²	0.2 (1.2)	0.1 (1.2)	0.0 (1.3)	-0.1 (1.4)	-0.2 (1.5)	-0.3 (1.6)	-0.6 (2.0)		<.001
Systolic blood pressure, mmHg	117.6 (11.9)	118.5 (11.5)	120.0 (11.1)	121.3 (10.7)	122.7 (10.2)	123.8 (9.8)	124.7 (8.9)		<.001
Diastolic blood pressure, mmHg	70.0 (8.5)	70.9 (8.3)	72.1 (8.1)	73.3 (7.9)	74.3 (7.6)	75.3 (7.5)	76.7 (7.4)		<.001
Total cholesterol, mmol/liter	5.1 (0.9)	5.2 (0.9)	5.3 (0.9)	5.4 (0.9)	5.4 (0.9)	5.5 (0.9)	5.5 (0.9)		<.001
HDL cholesterol, mmol/liter	1.7 (0.4)	1.6 (0.4)	1.5 (0.4)	1.4 (0.3)	1.4 (0.3)	1.3 (0.3)	1.3 (0.3)		<.001
Triacylglycerol, mmol/liter	1.0 (0.5)	1.2 (0.6)	1.3 (0.7)	1.5 (0.9)	1.6 (0.9)	1.8 (1.0)	1.9 (1.0)		<.001
Lipid medication use, %	1.7	1.5	2.3	2.9	3.3	3.1	3.2		0.367
Diabetic medication use, %	0.9	1.0	1.5	1.2	1.6	1.8	2.2		<.001
Blood glucose status, %									<.001
Normal	91.1	92.5	92.6	91.8	90.0	89.1	87.0		
Border	7.5	6.1	6.1	6.3	8.0	8.4	9.7		
Hyperglycemia	1.3	1.4	1.3	1.9	1.9	2.6	3.3		
Smoking status, %									<.001
Never	92.6	94.7	94.7	95.2	94.9	94.5	92.2		
Ex	0.6	0.5	0.7	0.5	0.6	0.9	0.9		
Current									
< 20 cigarettes/day	4.7	3.4	3.2	2.8	2.9	3.0	3.6		
≥ 20 cigarettes/day	2.1	1.5	1.5	1.5	1.6	1.6	3.3		
Alcohol intake, %									<.001
Never	90.9	88.7	89.0	89.6	90.2	90.7	90.7		
Sometimes	5.2	6.9	7.2	6.8	6.5	6.0	5.4		
Everyday									
< 66 g/day	3.9	4.2	3.7	3.5	3.3	3.1	3.4		
≥ 66 g/day	-	0.1	0.1	0.1	0.0	0.2	0.4		

HDL = high-density lipoprotein. The means (SD) are shown for continuous variables: age, fasting and non-fasting blood glucose, systolic and diastolic blood pressure, total cholesterol, HDL cholesterol, and triglycerides. SI conversion factors: blood glucose values were converted to mmol/L by multiply by 0.05551; cholesterol values were converted to mmol/L by multiply by 0.02586; triglycerides values were converted to mmol/L by multiply by 0.01129.

Table 2. The effect of hazard ratios on hypertension within each of seven BMI categories

	Baseline BMI categories (kg/cm ²)						
	< 19.0	19.0 - 20.9	21.0 - 22.9	23.0 - 24.9	25.0 - 26.9	27.0 - 29.9	> 30.0
Men, age 40-59 years (n = 8 540)							
No. of person-years	1 590	5 713	9 042	8 267	4 964	1 952	285
No. of hypertension cases	127	434	882	1 048	740	351	61
Incidence rates per 1 000 person-years	80	76	98	127	149	180	214
Age-adjusted HRs (95% CIs)	1.00	1.01 (0.83-1.23)	1.26 (1.05-1.52)	1.57 (1.31-1.89)	1.81 (1.50-2.19)	2.12 (1.73-2.60)	2.48 (1.83-3.36)
Model 1† HRs (95% CIs)	1.00	0.90 (0.74-1.09)	1.05 (0.87-1.27)	1.19 (0.98-1.44)	1.36 (1.16-1.66)	1.52 (1.23-1.89)	1.75 (1.28-2.40)
Model 2‡ HRs (95% CIs)	1.00	1.01 (0.59-1.73)	0.97 (0.58-1.65)	1.10 (0.64-1.88)	1.82 (1.04-3.16)	1.83 (1.00-3.35)	2.37 (0.91-6.17)
Model 3§ HRs (95% CIs)	1.00	0.95 (0.78-1.15)	1.13 (0.94-1.36)	1.24 (1.03-1.50)	1.42 (1.17-1.73)	1.60 (1.30-1.97)	1.77 (1.29-2.41)
Men, age 60-79 years (n = 9 796)							
No. of person-years	3 831	6 982	8 334	6 582	2 937	1 075	101
No. of hypertension cases	517	1 177	1 566	1 366	710	314	38
Incidence rates per 1 000 person-years	135	169	188	208	242	292	376
Age-adjusted HRs (95% CIs)	1.00	1.25 (1.13-1.38)	1.38 (1.25-1.52)	1.51 (1.36-1.67)	1.67 (1.49-1.87)	1.94 (1.69-2.24)	2.24 (1.61-3.11)
Model 1† HRs (95% CIs)	1.00	1.15 (1.03-1.27)	1.20 (1.09-1.33)	1.26 (1.13-1.41)	1.35 (1.20-1.53)	1.54 (1.32-1.78)	1.68 (1.20-2.35)
Model 2‡ HRs (95% CIs)	1.00	1.01 (0.71-1.44)	1.24 (0.88-1.75)	1.48 (1.03-2.13)	1.52 (0.98-2.37)	2.01 (1.11-3.64)	N/A
Model 3§ HRs (95% CIs)	1.00	1.11 (1.00-1.22)	1.17 (1.06-1.29)	1.26 (1.14-1.40)	1.34 (1.19-1.51)	1.48 (1.28-1.72)	1.68 (1.21-2.35)
Women, age 40-59 years (n = 32 133)							
No. of person-years	10 877	32 781	45 716	34 389	16 754	7 410	1 402
No. of hypertension cases	496	1 682	2 975	2 969	1 940	1 001	265
Incidence rates per 1 000 person-years	46	51	65	86	116	135	189
Age-adjusted HRs (95% CIs)	1.00	1.15 (1.04-1.27)	1.40 (1.27-1.54)	1.76 (1.60-1.94)	2.25 (2.04-2.49)	2.56 (2.30-2.86)	3.32 (2.86-3.86)
Model 1† HRs (95% CIs)	1.00	1.04 (0.94-1.15)	1.15 (1.05-1.27)	1.33 (1.20-1.46)	1.58 (1.43-1.75)	1.65 (1.48-1.84)	1.90 (1.63-2.22)
Model 2‡ HRs (95% CIs)	1.00	1.12 (0.90-1.14)	1.23 (0.99-1.53)	1.59 (1.28-1.98)	1.67 (1.32-2.11)	1.99 (1.52-2.60)	1.86 (1.16-2.98)
Model 3§ HRs (95% CIs)	1.00	0.98 (0.89-1.08)	1.12 (1.02-1.23)	1.27 (1.16-1.39)	1.47 (1.33-1.62)	1.55 (1.39-1.73)	1.91 (1.66-2.21)
Women, age 60-79 years (n = 17 736)							
No. of person-years	5 535	10 525	15 260	12 907	7 673	3 611	598
No. of hypertension cases	790	1 591	2 580	2 569	1 663	940	190
Incidence rates per 1 000 person-years	143	151	169	199	217	260	318
Age-adjusted HRs (95% CIs)	1.00	1.11 (1.02-1.21)	1.24 (1.14-1.34)	1.42 (1.31-1.54)	1.53 (1.40-1.66)	1.77 (1.61-1.95)	2.07 (1.77-2.43)
Model 1† HRs (95% CIs)	1.00	1.04 (0.95-1.13)	1.09 (1.01-1.19)	1.21 (1.11-1.31)	1.25 (1.15-1.37)	1.39 (1.26-1.54)	1.61 (1.37-1.90)
Model 2‡ HRs (95% CIs)	1.00	1.04 (0.80-1.35)	1.16 (0.90-1.49)	1.25 (0.96-1.62)	1.35 (1.02-1.80)	1.20 (0.84-1.70)	1.66 (0.82-3.38)
Model 3§ HRs (95% CIs)	1.00	1.06 (0.98-1.16)	1.10 (1.02-1.19)	1.22 (1.12-1.32)	1.29 (1.18-1.41)	1.40 (1.27-1.54)	1.72 (1.47-2.01)

Abbreviations: HRs, hazard ratios; CIs, confidence intervals; N/A, data was not available.


†Adjusted for age (years), systolic blood pressure (mmHg), diastolic blood pressure (mmHg), fasting status (yes or no), total cholesterol level (mmol/liter), high density lipoprotein cholesterol level (mmol/liter), log-transformed triglyceride level (mmol/liter), lipid medication use (yes or no), blood glucose status (normal, border, hyperglycemia), diabetic medication use (yes or no), smoking status (never, ex-, < 20 cigarettes/day, or ≥ 20 cigarettes/day), and alcohol intake (never, sometimes, < 66 g/day, or ≥ 66 g/day).

‡Addition of BMI (kg/m²), total cholesterol level (mmol/liter), high-density lipoprotein cholesterol level (mmol/liter), and log-transformed triglyceride level (mmol/liter) change values (the endpoint minus the baseline) to Model 1 for adjustment.

§Adjusted for time-dependent covariates: age (years), systolic blood pressure (mmHg), diastolic blood pressure (mmHg), fasting status (yes or no), total cholesterol level (mmol/liter), high density lipoprotein cholesterol level (mmol/liter), log-transformed triglyceride level (mmol/liter), lipid medication use (yes or no), blood glucose status (normal, border, hyperglycemia), diabetic medication use (yes or no), smoking status (never, ex-, < 20 cigarettes/day, or ≥ 20 cigarettes/day), and alcohol intake (never, sometimes, < 66 g/day, or ≥ 66 g/day).

Bold values showed statistical significance ($P < 0.05$).

Figure 1



No. of subjects	14 055	740	676	2 174
No. of hypertension cases	2 877	165	165	662